

The Relationship of Hamstrings and Quadriceps Strength to Anterior Cruciate Ligament Injury in Female Athletes

Gregory D. Myer, MS, CSCS,*†‡ Kevin R. Ford, MS,*†§ Kim D. Barber Foss, MS, ATC,*†
Chunyan Liu, MS,*¶ Todd G. Nick, PhD,*¶ and Timothy E. Hewett, PhD*†**

Objective: To determine the association of quadriceps and hamstrings strength to anterior cruciate ligament (ACL) injury risk in female athletes. The primary hypothesis was that there would be decreased knee flexor and increased knee extensor strength in female athletes who went on to ACL injured status (FACL) compared to uninjured female (FC) and male (MC) control subjects.

Study Design: Matched case control.

Setting: Institutional Biomechanics Laboratory.

Participants: Prospectively measured FACL (n = 22) female athletes who subsequently suffered confirmed noncontact ACL ruptures (16 during soccer and 6 during basketball play) were matched (1:4 ratio) to female controls (FC; n = 88) using limb (dominant or nondominant), pubertal status, sport, and nearest height and mass. In addition, male controls (MC) were matched (1:1 ratio) to FACL to serve as a secondary comparative control.

Assessment of Risk Factors: Isokinetic (concentric) knee extension/flexion strength (300 degrees/s).

Results: FACL subjects had decreased hamstrings strength compared to MC (15%; 95% CI, 1 to 27%; $P = 0.04$). FC were not different from MC in hamstrings strength. Conversely, FACL subjects did not differ compared to the MC in quadriceps strength, and the FC demonstrated decreased quadriceps strength relative to MC (10%; 95% CI, 3 to 18%; $P = 0.01$).

Conclusions: The results of this investigation indicate that female athletes who suffered ACL injury subsequent to strength testing had a combination of decreased hamstrings strength but not quadriceps

strength compared to males. In direct contrast, female athletes who did not go on to ACL injury had decreased quadriceps strength and similar hamstrings strength compared to matched male athletes.

Key Words: knee flexor to extensor ratio, isokinetic outcome measure, anterior cruciate ligament injury prediction, knee torques, knee co-contraction

(*Clin J Sport Med* 2009;19:3–8)

INTRODUCTION

Female athletes who participate in high-risk sports suffer anterior cruciate ligament (ACL) injury at a 4- to 6-fold greater rate than male athletes.¹ Lack of active neuromuscular control, as evidenced by increased knee abduction motion and torque,² and passive stability of the joint, as evidenced by increased joint laxity,³ may destabilize the knee and are predictive of increased ACL injury risk in female athletes. ACL injury likely occurs under conditions of high dynamic loading of the knee joint, when active muscular restraints do not adequately compensate for and adequately dampen joint loads.⁴ Decreased neuromuscular control of the joint may place stress on the passive ligament structures that exceed the failure strength of the ligament.^{5,6} Neuromuscular control of high-load movements is required to maintain dynamic knee stability during landing and pivoting.^{5,7,8} Hamstrings and quadriceps co-contraction may provide dynamic joint stabilization and potentially protect the knee during sports-related tasks.^{5,8–10} Male athletes demonstrate increased relative strength^{11–13} and relative recruitment^{13,14} of the hamstrings compared to quadriceps during dynamic knee-loading tasks. Ford and colleagues reported that female athletes use increased quadriceps activation without matched increases in hamstrings activation when they performed drop landings with incrementally increased drop height intensity.¹⁵ Decreased hamstrings relative to quadriceps strength and recruitment is implicated as a potential mechanism for increased lower extremity injuries.^{10,15–18} Deficits in relative hamstrings strength and recruitment may also contribute to increased ACL injury risk in female athletes.

The purpose of the current investigation was to determine the association of quadriceps and hamstrings isokinetic strength to ACL injury risk in female athletes. The primary hypothesis was that decreased knee flexor and increased knee extensor strength would be observed in female

Submitted for publication June 18, 2008; accepted October 2, 2008.

From *Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio; †Sports Medicine Biodynamics Center and Human Performance Laboratory, Cincinnati, Ohio; ‡Rocky Mountain University of Health Professions, Provo, Utah; §Department of Kinesiology and Health Promotion, University of Kentucky, Lexington, Kentucky; ¶Center for Epidemiology and Biostatistics, Cincinnati, Ohio; and **Departments of Pediatrics, Orthopaedic Surgery, College of Medicine and the Departments of Biomedical Engineering and Rehabilitation Sciences, University of Cincinnati, Cincinnati, Ohio.

This work was supported by National Institutes of Health Grant R01-AR049735 and R01-AR055563.

The authors state that they have no financial interest in the products mentioned within this article.

Reprints: Gregory D. Myer, MS, CSCS, Cincinnati Children's Hospital, 3333 Burnet Avenue, MLC 10001, Cincinnati, OH 45229 (e-mail: greg.myer@chmcc.org).

Copyright © 2009 by Lippincott Williams & Wilkins

athletes who would subsequently suffer ACL injury (FACL) compared to uninjured female control (FC) and male control (MC) subjects.

METHODS

Subjects

A matched case control study design ($n = 132$ total, 22 FACL, 110 controls) was used to sample female and male high school and collegiate soccer and basketball players who were prospectively screened (2002 to 2007; $n = 1692$ potential samples) for hamstrings and quadriceps strength measures before their recorded ACL injury. FACL ($n = 22$) subsequently suffered arthroscopically confirmed noncontact ACL ruptures (16 during soccer and 6 during basketball play). FACL injuries occurred in a range of 1 to 24 months from date of testing. Each FACL case was matched to 4 FC ($n = 88$) using limb (dominant or nondominant), pubertal status, sport, and nearest height and mass. In addition, MC were also matched using limb, pubertal status, and nearest height and mass matched (1:1 ratio) to FACL to serve as a secondary comparative control.

Informed written consent was obtained from all subjects and their parents and was approved by the Institutional Review Board. After the informed consent was obtained, height and mass were measured and recorded. Isokinetic strength and vertical jump height measures were recorded during the laboratory evaluation during each athlete's visit for their prospective screening. Included within the consent form was a detailed medical history and previous sport participation questionnaire.

Maturational Status

The modified Pubertal Maturational Observational Scale (PMOS) has been shown to provide reliable assessment of pubertal status, and 1 rater was used to classify each subject in the current study into 1 of 3 pubertal categories (prepubertal, pubertal, and postpubertal).^{19–23}

Vertical Jump Height Testing

Figure 1 presents vertical jump measurement on the MX1 vertical jump trainer (MXP Sports, Reading, Pennsylvania). A previously published report demonstrated that countermovement vertical jump testing has a test-retest reliability of $R = 0.993$.²⁴

Dynamic Strength

Isokinetic knee extension/flexion (concentric/concentric muscle action) strength was collected with each subject seated on the dynamometer with trunk perpendicular to floor, the hip flexed to 90 degrees, and the knee flexed to 90 degrees. Before each data collection set, a warm-up set consisting of 5 submaximal knee flexion/extensions for each leg at 300 degrees/s was performed. The test session consisted of 10 knee isokinetic extension/flexion (100 to 0 degrees range of motion) repetitions for each leg at 300 degrees/s (Figure 2). Peak flexion and extension torques were recorded (ft·lbs). Isokinetic concentric strength measures of hamstrings and quadriceps strength demonstrate excellent reliability.²⁵



FIGURE 1. (A) Example of reach testing. (B) Example of maximum effort countermovement vertical jump testing.

Statistical Method

Descriptive statistics were calculated to characterize the study population. As a result of the skewed nature of the data, medians and interquartile ranges (the difference between 25th and 75th percentiles) were reported. Natural log transformations were taken on all response measures and used for subsequent modeling. Logs were found to be appropriate on the basis of the Box-Cox transformation computed using PROC TRANSREG in SAS (SAS Institute, Cary, North Carolina). To account for the matching strategy, a linear mixed model using PROC MIXED in SAS was fitted on each response to compare case-control differences. The matching variable was treated as a random effect. To take into account the difference in individual physical capabilities, vertical jump



FIGURE 2. Example of the isokinetic knee flexion/extension testing.

height was added to each model as a control variable to explain part of the variability in response. To relate to the original scale of the data, a back-transform (antilog) was used on the mean log difference to compute the geometric mean of a ratio defined as the ratio of one group compared to a reference group.

RESULTS

The matched case control study design yielded 110 female and 22 male middle school, high school, and collegiate soccer and basketball players who were all prospectively screened for strength measures before the recorded ACL injury for the FACL subjects. The FACL sample was similar in height (FACL, 165.6 ± 6.5 cm; FC, 164.9 ± 5.5 cm; MC, 167.2 ± 5.7 cm; *P* = 0.25) and mass (FACL, 58.8 ± 7.3; FC, 58.2 ± 7.1; MC, 60.2 ± 7.7 kg; *P* = 0.51) to their matched male and female controls. Nineteen of the FACL were categorized as postpubertal, and three were categorized as pubertal at time of assessment.

Figure 3 presents the median values for hamstrings to quadriceps strength ratio (H/Q), and the interquartile ranges of the recorded hamstrings peak torque, quadriceps peak torque, and vertical jump height for each group are presented in Table 1. These were unadjusted measurements and did not account for matching. The comparisons in each response between the matched groups are summarized in Figure 4. Because the geometric means were used for the log-transformed data, the results can be interpreted in terms of percent change.

Female athletes who subsequently suffered ACL injury had decreased hamstrings strength, but not decreased similar quadriceps strength, compared to matched control males. Conversely, female athletes who did not go on to ACL injury had decreased quadriceps strength, but not decreased




FIGURE 3. Median hamstrings to quadriceps ratio for each of the matched comparison groups. FACL, female ACL injured; FC, female control; MC, male control.

TABLE 1. Quartiles (25th, 50th, 75th Percentiles) of Response Variables (Quadriceps and Hamstrings Strength) and Covariate (Vertical Jump) by Study Group

Response	FACL (n = 22)	FC (n = 88)	MC (n = 22)
Quadriceps (ft-lbs)	(51.9, 56.6, 65.6)	(51.1, 56.4, 61.5)	(61.0, 67.6, 73.2)
Hamstrings (ft-lbs)	(29.2, 34.5, 37.9)	(27.9, 33.6, 39.4)	(34.8, 38.2, 42.5)
Vertical jump (cm)	(14.0, 15.8, 17.0)	(14.0, 15.5, 17.0)	(17.0, 19.7, 21.0)

FACL, female ACL injured; FC, female control; MC, male control.

hamstrings strength, compared to matched male athletes. Although the median measurements in quadriceps and hamstrings strength, were similar in FACL and FC groups, Figure 4 shows that FACL subjects had 15% decreased hamstrings strength compared to MC (95% CI, 1 to 27%; *P* = 0.04), and FC were not different than MC in hamstrings strength (*P* = 0.08) after adjusting for vertical jump height and taking into account the random effects arising from the matching. Conversely, the FC demonstrated a 10% decrease in quadriceps strength relative to MC (95% CI, 3 to 18%; *P* = 0.01). FACL subjects did not differ compared to the MC in quadriceps strength (*P* = 0.14).

DISCUSSION

Previously published critiques of similar epidemiological designs to determine risk factors for ACL injury have expressed concern about the confounding effects of sport, height, and mass.^{2,26} This investigation incorporated an approach to control for potential confounding variance among individual sport, maturational status, and individual anthropometrics by using matched case subjects with subjects of the same sport, pubertal status, and nearest height and mass to serve as uninjured control subjects. In addition, this study used a novel approach to compare matched male athletes to control




FIGURE 4. Percent increase in responses and 95% CI comparing study group to reference group. FACL, female ACL injured; FC, female control; MC, male control. **P* < 0.05.

for potential confounding variance in whole body power and maturational status between investigational and control subjects. Cumulatively, the control measures in this investigation were employed to provide a more robust interpretation of isolated strength mechanisms and their relationship to ACL injury in female athletes. The results of the current investigation indicate that, when controlling for whole body power, FACL demonstrate similar relative quadriceps strength, but decreased hamstrings strength, compared to uninjured male athletes. This finding indicates that decreased relative hamstrings strength and recruitment may be a potential contributing mechanism to ACL injury in high-risk female athletes.

A previous study showed that male and female relative hamstrings to quadriceps strength profiles diverge significantly during and after puberty.²¹ Isokinetic dynamometer measurements show that male athletes demonstrate significantly greater hamstrings peak torques with increasing maturity, while peak hamstrings torque remain stable with increasing maturational stage in female athletes.²¹ In addition, sex differences in isokinetic H/Q ratios are not detected in postpubertal athletes at slower angular velocities. However, at high knee flexion/extension angular velocities approaching those that occur during sports activities, significant gender differences become evident in H/Q ratio.¹¹ Unlike their male counterparts, female athletes do not increase hamstrings to quadriceps torque ratios at velocities that approach those of functional activities.¹¹ Thus, it appears that decreased hamstrings strength and H/Q ratios of female athletes relative to males may be related to the development of neuromuscular imbalances associated with the onset of maturation.^{12,21,27} These neuromuscular imbalances may increase ACL injury risk in pubertal and postpubertal female athletes, as was indicated by the current results with all FACL subjects either pubertal or postpubertal at the time of injury.^{2,21,28}

Decreased hamstrings strength relative to the quadriceps is implicated as a potential mechanism for increased lower extremity injuries,^{8,15-17} and potentially ACL injury risk, in female athletes.¹⁸ Joint stability through hamstrings and quadriceps co-contraction may be necessary when the joint experiences high quadriceps activation or when the passive structures are compromised.^{29,30} Withrow and colleagues reported that increased hamstrings force during the flexion phase of simulated jump landings greatly decreased relative strain on the ACL.¹⁰ Another proposed theory related to neuromuscular imbalances and increased ACL injury risk in females is the relatively low knee flexor to extensor recruitment, which is reflective of a closed chain dynamic H/Q.^{13,14} For example, hamstrings activation can decrease the load on the passive restraints of the knee,³¹ increase the knee joint compression force, and stabilize the knee from external varus/valgus load.³² Hewett et al¹³ reported that male athletes demonstrated knee flexor moments measured using inverse dynamics that were 3-fold higher than female athletes when decelerating from a landing. This group of female athletes also demonstrated decreased isokinetically measured H/Q ratio and increased knee abduction (valgus) moments compared to male subjects. The increased knee valgus moment significantly correlated to the peak impact forces during the maneuver and is purported to increase ACL injury risk in

female athletes.^{2,13} Ford et al¹⁵ reported that female athletes showed an absence of matched increased hamstrings muscle activation relative to quadriceps and overall low amplitude during the landing phase of a jump. This tendency of female athletes to preferentially activate the quadriceps relative to the hamstrings during high-demand activities may limit their ability to maintain dynamic knee control during high-risk maneuvers.

This quadriceps dominance, or decreased recruitment of hamstrings relative to the quadriceps, has also been observed in elite female collegiate athletes.³³ Female athletes reacted to a forward translation of the tibia primarily with a muscular activation of the quadriceps muscles, whereas male athletes relied on their hamstrings muscles to counteract the anterior tibial displacement.³³ Malinzak et al³⁴ examined male and female athletes during more sport specific tasks of running, cross-cutting and side-cutting tasks. They found that female athletes used less knee flexion, increased quadriceps activation, and decreased hamstrings activation compared to male athletes when performing the 3 athletic tasks.³⁴ In addition, Sigward and Powers³⁵ examined a side-step cutting maneuver and found that skilled female soccer players had less co-contraction than novice players. These findings indicate that a movement pattern may be learned, which may elevate possible injury risk in skilled players. Recently, Lawrence et al³⁶ reported that female athletes in the decreased hip and hamstrings strength group demonstrated increased relative knee extensor activity, ground reaction force, and coronal plane knee load when landing on a single leg. Cumulatively, this evidence indicates that decreased relative hamstrings strength may underlie high-risk biomechanics that contribute to ACL injury in female athletes.^{2,13,30,36}

The tendency to land with a straighter knee during high-intensity tasks could be exacerbated by early quadriceps or delayed hamstrings activation during weight-bearing stance.³⁷ Chappell et al³⁸ concluded that the increased anterior shear force demonstrated by the female athletes was potentially due to the combination of increased quadriceps force, decreased hamstrings force, and decreased knee flexion. A sagittal plane position of the knee near full extension when landing or cutting is commonly observed in video analysis of ACL injuries in female athletes.^{39,40} In addition, a prospective study indicated that female athletes who subsequently sustained ACL injuries demonstrated significantly less (10.5 degrees) knee flexion when performing a drop vertical jump than those who did not sustain injury.² At low knee flexion angles (0 to 30 degrees of knee flexion), quadriceps contractions pull the tibia forward and increase stress on the ACL, especially without balanced knee flexor (hamstrings and gastrocnemius) co-contraction to decrease strain on the ligament.^{29,38,41} Performing tasks with relatively low knee flexion angles (0 to 30 degrees of knee flexion) in conjunction with relatively large quadriceps contractions may elicit anterior tibial shear forces that are large enough to rupture the ACL.^{38,42} Increased hamstrings strength and recruitment may decrease ACL injury risk via a mechanism of increased relative hamstrings co-contraction that may lead to increased knee flexion and reduced knee abduction and anterior tibial shear during dynamic tasks.^{2,13,29,39,40,43-46} The FACL subjects in the current study with decreased relative hamstrings strength may have

also demonstrated decreased hamstrings recruitment during dynamic tasks, which may have contributed to their injury mechanism; however, muscle activation patterns were not examined in the current study, and these neuromuscular parameters cannot be directly determined.

Comprehensive neuromuscular training that combines resistance training with either plyometric training or dynamic balance training can increase knee flexion and decrease dangerous coronal plane knee loads during dynamic tasks, with concomitant increases in relative hamstrings strength in female athletes.^{13,45-47} However, isolated resistance training protocols appear to reduce relative hamstrings to quadriceps ratio and are not effective to alter potential high-risk sagittal or coronal plane biomechanics in female athletes.^{48,49} The increased ability to decelerate from a landing and to control dynamic knee valgus may be related to the decreased imbalance in the hamstrings to quadriceps strength and recruitment that was observed in the female athletes before training.⁵⁰ The increased balance in strength and recruitment of the hamstrings and gastrocnemius musculature relative to quadriceps may be a mechanism that protects the knee ligaments in male athletes and trained female athletes.^{2,13,43} Adequate co-contraction of the knee flexors may help balance active contraction of the quadriceps that can compress the joint and assist in the control of high knee abduction torques or valgus collapse. Appropriate neuromuscular control may prevent the critical loading necessary to rupture the ACL during maneuvers that place the athlete at risk for an injury. Female athletes, with decreased ability to adequately balance muscular recruitment through positions of high joint loading, may increase their risk of subsequent ligament failure.^{2,13,29,39,40,43-46} Therefore, dynamic knee stability during multiplanar movements at the knee may be contingent on hamstrings strength and coactivation to resist anterior tibial translation and knee abduction resulting from excessive quadriceps contraction.^{13,44,51} On the basis of the results of the current report, increased relative strength and recruitment of the hamstrings musculature achieved through neuromuscular training may be a potential mechanism for successful reduction of ACL injury in female athletes who lack dynamic knee stability.^{2,13,43,45,46}

Study Limitations

ACL injury in female athletes likely has a multifactorial etiology, with several potential elements that determine injury mechanism. Specifically, increased knee abduction motion and torque² and increased knee joint laxity³ predict increased ACL injury risk in female athletes. These epidemiologic data demonstrate that biomechanical factors may contribute in isolation or combination with other factors such as anatomical, hormonal, and potentially psychological parameters to increase relative ACL injury rates in female athletes. Although this investigation made several efforts to provide the most appropriate control available to sample, it is plausible that contributing and confounding variables were not controlled for in the current study design.

In addition, the muscular strength contributions that occur during ACL injury occur at a high rate of knee joint loading in closed chain situations, with the quadriceps most

likely acting with eccentric muscle action with the hip angles more extended than the current test protocol.⁵² While the current protocol did measure the strength at relatively high speeds (300 degrees/s) the difference in muscle action (concentric versus eccentric) of the quadriceps during open chain measurements may limit the applicability of our findings to conditions that occur during actual ACL injury and limit our potential to determine significant group differences in H/Q measures.

However, the significant effects of group differences in relative FACL hamstrings strength were observed in this study compared to carefully matched males. These differences were not evident in the uninjured FC compared to the MC. The observed differences were of a magnitude outside of the protocol's measurement error and should be considered a potential contributing mechanism to increased ACL injury risk in female athletes.

CONCLUSIONS

The results of this investigation indicate that female athletes who went on to suffer ACL injury had a combination of decreased hamstrings strength, but not decreased similar quadriceps strength, compared to male athletes. In direct contrast, female athletes who did not go on to ACL injury had decreased quadriceps strength, but not decreased hamstrings strength, compared to matched male athletes. Female athletes who demonstrate the combination of decreased relative hamstrings and high relative quadriceps strength may be at increased risk for ACL injury. Preseason screening programs that monitor hamstrings and quadriceps strength, especially relative to comparable values in male athletes, may be warranted to identify female athletes with potential deficits. Targeted neuromuscular interventions that increase relative hamstrings muscle strength and recruitment may decrease injury risk and potentially increase performance in this population.

ACKNOWLEDGMENTS

We thank Jurdan Mendigutxia for his critical input into the manuscript content. We also thank Boone County School District Kentucky, especially School Superintendents Randy Poe and Dr. Brian Blavatt, for participation in this study.

REFERENCES

1. Arendt E, Dick R. Knee injury patterns among men and women in collegiate basketball and soccer. NCAA data and review of literature. *Am J Sports Med.* 1995;23:694-701.
2. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005;33:492-501.
3. Myer GD, Ford KR, Paterno MV, et al. The effects of generalized joint laxity on risk of anterior cruciate ligament injury in young female athletes. *Am J Sports Med.* 2008;36:1073-1080.
4. Beynon BD, Fleming BC. Anterior cruciate ligament strain in-vivo: a review of previous work. *J Biomech.* Jun 1998;31:519-525.
5. Li G, Rudy TW, Sakane M, et al. The importance of quadriceps and hamstring muscle loading on knee kinematics and in-situ forces in the ACL. *J Biomech.* Apr 1999;32:395-400.

6. Markolf KL, Graff-Redford A, Amstutz HC. In vivo knee stability: a quantitative assessment using an instrumented clinical testing apparatus. *J Bone Joint Surg Am.* 1978;60A:664–674.
7. Besier TF, Lloyd DG, Cochrane JL, et al. External loading of the knee joint during running and cutting maneuvers. *Med Sci Sports Exerc.* 2001;33:1168–1175.
8. Ford KR, van den Bogert AJ, Myer GD, et al. The effects of age and skill level on knee musculature co-contraction during functional activities: a systematic review. *Br J Sports Med.* 2008;42:561–566.
9. Renstrom P, Arms SW, Stanwyck TS, et al. Strain within the anterior cruciate ligament during hamstring and quadriceps activity. *Am J Sports Med.* 1986;14:83–87.
10. Withrow TJ, Huston LJ, Wojtys EM, et al. Effect of varying hamstring tension on anterior cruciate ligament strain during in vitro impulsive knee flexion and compression loading. *J Bone Joint Surg Am.* 2008;90:815–823.
11. Hewett TE, Myer GD, Zazulak BT. Hamstrings to quadriceps peak torque ratios diverge between sexes with increasing isokinetic angular velocity. *J Sci Med Sport.* 2008;11:452–459.
12. Holm I, Vollestad N. Significant effect of gender on hamstring-to-quadriceps strength ratio and static balance in prepubescent children from 7 to 12 years of age. *Am J Sports Med.* 2008;36:2007–2013.
13. Hewett TE, Stroupe AL, Nance TA, et al. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996;24:765–773.
14. Malinzak RA, Colby SM, Kirkendall DT, et al. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech.* 2001;16:438–445.
15. Ford KR, Myer GD, Schmitt LC, et al. Effect of drop height on lower extremity biomechanical measures in female athletes. *Medicine & Science in Sports & Exercise.* 2008;40:S80.
16. Knapik JJ, Bauman CL, Jones BH, et al. Preseason strength and flexibility imbalances associated with athletic injuries in female collegiate athletes. *Am J Sports Med.* 1991;19:76–81.
17. Soderman K, Alfredson H, Pietila T, et al. Risk factors for leg injuries in female soccer players: a prospective investigation during one out-door season. *Knee Surg Sports Traumatol Arthrosc.* 2001;9:313–321.
18. Myer GD, Ford KR, Hewett TE. Rationale and clinical techniques for anterior cruciate ligament injury prevention among female athletes. *J Athl Train.* 2004;39:352–364.
19. Quatman CE, Ford KR, Myer GD, et al. The effects of gender and pubertal status on generalized joint laxity in young athletes. *J Sci Med Sport.* 2008;11:257–263.
20. Quatman CE, Ford KR, Myer GD, et al. Maturation leads to gender differences in landing force and vertical jump performance: a longitudinal study. *Am J Sports Med.* 2006;34:806–813.
21. Hewett TE, Myer GD, Ford KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *J Bone Joint Surg Am.* 2004;86-A:1601–1608.
22. Davies PL, Rose JD. Motor skills of typically developing adolescents: awkwardness or improvement? *Phys Occup Ther Pediatr.* 2000;20:19–42.
23. Davies PS. *Assessment of cognitive development in adolescents by means of neuropsychological tasks.* [Dissertation]. Laramie: Department of Psychology, University of Wyoming; 1995.
24. Stockbrugger BA, Haennel RG. Validity and reliability of a medicine ball explosive power test. *J Strength Cond Res.* 2001;15:431–438.
25. Bandy WD, McLaughlin S. Intramachine and intermachine reliability for selected dynamic muscle performance tests. *J Orthop Sports Phys Ther.* 1993;18:609–613.
26. Orchard JW. Potential to reduce the risk of noncontact anterior cruciate ligament (ACL) injuries [letter]. *Am J Sports Med.* 2005;33:1930; author reply, 1930–1931.
27. Ahmad CS, Clark AM, Heilmann N, et al. Effect of gender and maturity on quadriceps-to-hamstring strength ratio and anterior cruciate ligament laxity. *Am J Sports Med.* 2006;34:370–374.
28. Shea KG, Pfeiffer R, Wang JH, et al. Anterior cruciate ligament injury in pediatric and adolescent soccer players: an analysis of insurance data. *J Pediatr Orthop.* 2004;24:623–628.
29. Sell TC, Ferris CM, Abt JP, et al. Predictors of proximal tibia anterior shear force during a vertical stop-jump. *J Orthop Res.* 2007;25:1589–1597.
30. Withrow TJ, Huston LJ, Wojtys EM, et al. The relationship between quadriceps muscle force, knee flexion, and anterior cruciate ligament strain in an in vitro simulated jump landing. *Am J Sports Med.* 2006;34:269–274.
31. MacWilliams BA, Wilson DR, DesJardins JD, et al. Hamstrings cocontraction reduces internal rotation, anterior translation, and anterior cruciate ligament load in weight-bearing flexion. *J Orthop Res.* 1999;17:817–822.
32. Lloyd DG, Buchanan TS. Strategies of muscular support of varus and valgus isometric loads at the human knee. *J Biomech.* 2001;34:1257–1267.
33. Huston LJ, Wojtys EM. Neuromuscular performance characteristics in elite female athletes. *Am J Sports Med.* 1996;24:427–436.
34. Malinzak RA, Colby SM, Kirkendall DT, et al. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech (Bristol, Avon).* 2001;16:438–445.
35. Sigward S, Powers CM. The influence of experience on knee mechanics during side-step cutting in females. *Clin Biomech (Bristol, Avon).* 2006;21:740–747.
36. Lawrence RK, 3rd, Kernozek TW, Miller EJ, et al. Influences of hip external rotation strength on knee mechanics during single-leg drop landings in females. *Clin Biomech (Bristol, Avon).* 2008;23:806–813.
37. Shultz SJ, Perrin DH, Adams MJ, et al. Neuromuscular response characteristics in men and women after knee perturbation in a single-leg, weight-bearing stance. *J Athl Train.* 2001;36:37–43.
38. Chappell JD, Yu B, Kirkendall DT, Garrett WE. A comparison of knee kinetics between male and female recreational athletes in stop-jump tasks. *Am J Sports Med.* 2002;30:261–267.
39. Olsen OE, Myklebust G, Engebretsen L, et al. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med.* 2004;32:1002–1012.
40. Boden BP, Dean GS, Feagin JA, et al. Mechanisms of anterior cruciate ligament injury. *Orthopedics.* 2000;23:573–578.
41. Cowling EJ, Steele JR. The effect of upper-limb motion on lower-limb muscle synchrony. Implications for anterior cruciate ligament injury. *J Bone Joint Surg Am.* 2001;83-A:35–41.
42. DeMorat G, Weinhold P, Blackburn T, et al. Aggressive quadriceps loading can induce noncontact anterior cruciate ligament injury. *Am J Sports Med.* 2004;32:477–483.
43. Hewett TE, Lindenfeld TN, Riccobene JV, et al. The effect of neuromuscular training on the incidence of knee injury in female athletes. A prospective study. *Am J Sports Med.* 1999;27:699–706.
44. White KK, Lee SS, Cutuk A, et al. EMG power spectra of intercollegiate athletes and anterior cruciate ligament injury risk in females. *Med Sci Sports Exerc.* 2003;35:371–376.
45. Myer GD, Ford KR, McLean SG, et al. The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics. *Am J Sports Med.* 2006;34:490–498.
46. Myer GD, Ford KR, Brent JL, et al. The effects of plyometric versus dynamic balance training on power, balance and landing force in female athletes. *J Strength Cond Res.* 2006;20:345–353.
47. Myer GD, Ford KR, Palumbo JP, et al. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *J Strength Cond Res.* 2005;19:51–60.
48. McGinn PA, Mattacola CG, Malone TR, et al. Strength training for 6-weeks does not significantly alter landing mechanics of female collegiate basketball athletes. *J Orthop Sports Phys Ther.* 2007;37:A24.
49. Herman DC, Weinhold PS, Guskiewicz KM, et al. The effects of strength training on the lower extremity biomechanics of female recreational athletes during a stop-jump task. *Am J Sports Med.* 2008;36:733–740.
50. Hewett TE, Stroupe AL, Nance TA, et al. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996;24:765–773.
51. Draganich LF, Vahey JW. An in vitro study of anterior cruciate ligament strain induced by quadriceps and hamstrings forces. *J Orthop Res.* 1990;8:57–63.
52. Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med.* 2007;35:359–367.